

REDESCRIPTION OF A NORTH AMERICAN FRESHWATER PLANARIAN,
SEIDLIA REMOTA (SMITH, 1988), WITH TAXONOMIC NOTES
ON *SEIDLIA* AND *POLYCELIS* SPECIES FROM THE FAR EAST
AND CENTRAL ASIA
(TURBELLARIA, SERIATA, TRICLADIDA, PALUDICOLA)¹⁾

by

MASAHARU KAWAKATSU and ROBERT W. MITCHELL

INTRODUCTION

A Eurasian genus *Polycelis* EHRENBERG, 1831, in the family Planariidae STIMPSON, 1857, of freshwater planarian has over 30 species and several subspecies in Europe, Central Asia, the Far East, and the United States. The genus is, however, not homogeneous according to details of the genital anatomy and histology (especially in the anatomy of the copulatory apparatus).

The Russian taxonomists (ZABUSOV, 1911, 1916, ZABUSOVA, 1929, 1936; LIVANOV & ZABUSOVA, 1940) proposed the 5 generic classification system: *Polycelis* EHRENBERG, 1831; *Sorocelides* ZABUSOVA, 1929; *Polycelidia* ZABUSOVA, 1936; *Seidlina* ZABUSOV, 1911; *Ijimia* BERGENDAL, 1890. KENK (1930, 1953, 1972, 1973, 1974, 1989) proposed the 3 subgeneric classification system: *Polycelis* (syn. *Sorocelides* and *Polycelidia*), *Seidlina* and *Ijimia* (cf. KAWAKATSU, OKI, TAMURA, TAKI, YAMAMOTO, NISHINO, TIMOSHKIN, KUZNEDELOV & SLUYS, 1996, p. 16).

The main subgeneric characters in KENK's (*op. cit.*) system are: 1) *Polycelis* (lacking adenodactyls and lacking excessive development of the muscle coat of the male genital antrum); 2) *Seidlina* (having an extraordinarily thick muscle zone surrounding the male genital antrum and without adenodactyls); 3) *Ijimia* (possessing solid adenodactyls and without thick muscle zone of the male genital antrum).

KAWAKATSU & MITCHELL (1995) considered that the subgenus *Seidlina* should be recognized at the rank of genus to which it was elevated by ZABUSOVA in 1936.

According to this new taxonomic system, the genus *Polycelis* (*s.l.*) is separable as follows: the genera *Polycelis* EHRENBERG, 1831 (consists of the subgenera *Polycelis* EHRENBERG, 1831, and *Ijimia* BERGENDAL, 1890) and *Seidlina* ZABUSOV, 1911.

The species of the subgenus *Polycelis* show a wide geographical distribution in Europe, Central Asia, the Far East, and the United States. The species of the subgenus *Ijimia* is distributed only in Europe (a single species is distributed in New Guinea). The *Seidlina* species are distributed in Central Asia, the Far East and United States.

The first *Seidlina* species reported from the United States was *Seidlina sierrensis* (KENK, 1973) (*olim Polycelis* (*Seidlina*) *sierrensis*) from California and Nevada. In 1988, a new species, "*Polycelis remota* SMITH, 1988", was described from a spring in Western Massachusetts. KAWAKATSU felt that misunderstandings of penial anatomy appeared in the original description (Biol. Bull., 175, fig. 2 on p. 248; parts of the organ represented as PV, ED, PB, and PP are incorrectly interpreted). We studied newly-prepared serial sections of "*Pol. remota*" from the type locality (obtained by the courtesy of Dr. D. G. SMITH). Microscopic examination revealed that the species is the second *Seidlina* species occurring in the United States.

In the present paper, a taxonomic redescription of *Seidlina remota* (SMITH, 1988) will be given first. Following, taxonomic notes on *Seidlina* and *Polycelis* species from the Far East and Central Asia will be given because several new species (including synonyms) in these genera have

1) Parts of this paper were given at the 66th Annual Meeting of the Zoological Society of Japan held in Hachijōji, Tōkyō, Japan (September 15-17, 1995; cf. KAWAKATSU & MITCHELL, 1995) and the VIIIth International Symposium on the Biology of Turbellaria held in Brisbane, Australia (August 19-23, 1996; cf. KAWAKATSU & TIMOSHKIN, 1996, 1998; see also KAWAKATSU, OKI, TAMURA, TAKAI, YAMAMOTO, NISHINO, TIMOSHKIN, KUZNEDELOV & SLUYS, 1996).

been reported after publication of KENK's "Index of the Genera and Species of the Freshwater Triclad (Turbellaria) of the World (1974)."

MATERIAL AND METHODS

The following samples of "*Polycelis remota*" were examined.

KAWAKATSU's Specimen Lot No. 1985. About 20 sexual and asexual specimens collected from the type locality: a spring in Sunderland, Franklin County, Massachusetts, U. S. A. (see SMITH, 1988, p. 246); fixed with FAA and then presently in 3% formalin solution. Coll. Dr. D. G. SMITH; November 7, 1989.

Serial sections (7-8 micrometers) of 13 specimens stained with Delafield's hematoxylin and erythrosin were prepared by KAWAKATSU. Whole mounts of 2 specimens (a single specimen without staining and another specimen stained with erythrosin) were also prepared for stereomicroscopic examination.

REDESCRIPTION OF "*POLYCELIS REMOTA* SMITH, 1988"

Order TRICLADIDA

Suborder PALUDICOLA or PROBURSALIA

Family PLANARIIDAE STIMPSON, 1857

Genus *Seidlia* ZABUSOV, 1911

Seidlia remota (SMITH, 1988)

Principal literature.

Polycelis remota SMITH, 1988, in Biol. Bull., 175: 246-252, figs. 1-3. Original description.

Seidlia remota in KAWAKATSU & MITCHELL, 1995, Zool. Sci., Tôkyô, 12-Suppl.: 32. Taxonomic remarks.

External features. External appearance of live specimens of this species with sketch figures is described by SMITH (1988, p. 247, fig. 1 A-C; see Fig. 1 A-C in the present paper). Preserved sexually mature specimens measure 9 to 12mm long and 1.5-2.5 mm in width. The head is gently convex with moderately-developed, prominent auricles (Figs. 2 A-C, 3 A and B). Behind the head, the "neck" is conspicuous.

The ground color of the dorsal side of the body is pale redish brown. The ventral side is a lighter hue. A pharynx is inserted somewhat behind the middle of the body. The copulatory organs occupy more than the anterior one-third of the post-pharyngeal region, so that the genital pore opens to a slightly anterior level (Figs. 2 A-C, 3).

SMITH (1988, p. 247, fig. 1 D; see Fig. 1 D in the present paper) described a presence of "a single exterior sucker-like organ situated always on the right ventral side of the animal, anterior and separate from the gonopore and its associated atrial cavities." However, no such structure was found in our samples examined (this is also confirmed by the examination of our serial sections).

The species has many small eyes near the anterior and the lateral margins of the head (Figs. 2A, 3 A and B; see also Fig. 1 A-C). The eyes in a curved band of two to three rows are absent at the top of the frontal margin and the basal portions of the auricles; behind the level of auricles, the band narrows to a single row that terminating at the anterior one-third level of the prepharyngeal region. The variations of number of eyes are shown in Table 1.

Judging from our data, the general arrangement of eyes in *S. remota* is different from that of North American *Seidlia sierrensis* KENK, 1973 (cf. KENK, 1973, p. 9, fig. 6 C). It is also different from that of 3 Japanese *Seidlia* species: *Seidlia auriculata* IJIMA et KABURAKI, 1916 (cf. KAWAKATSU, TESHIOGI & ISHIDA, 1980, p. 132, fig. 1); *Seidlia akkeshi* (cf. ICHIKAWA & KAWAKATSU, 1963, pp. 1-3, figs. 1 A-E and 2 A-F); *Seidlia schmidtii* (ZABUSOV, 1916) (cf. KAWAKATSU, 1964, pp. 176-177, figs. 1 A-C and 2 A-F; KAWAKATSU & YAMADA, 1966, p. 3, fig. 1 a-h).

Internal features. The pharynx has a structure typical of the genus *Seidlia* in the family Planariidae. The anterior intestinal trunk has about 10 short lateral branches; each posterior trunk has 12 to 15 short branches. In large specimens, two medial branches are united at the middle level of the copulatory apparatus (between the penis and the genital pore on the ventral side) and from a well-developed intestinal anastomosis. It is a wide canal lined with a very high, highly glandular epithelium. The muscular coating of the canal consists chiefly of a thick

Table 1. Variation of the number of eyes in *Seidlia remota* from the type locality
(KAWAKATSU's Specimen Lot No. 1985).

The body length was measured from the preserved specimens.

No. of specimens examined, their sexual state in parentheses	Body length \times width (mm)	No. of eyes		
		Minimum on one side	Maximum on one side	Total no. of eyes
15(13, sexual 2, asexual	8-11 \times 1.5-2	28	42	57-82
	4-5 \times 1	16	35	33-65




Fig. 1. "*Polycelis remota* SMITH, 1988", external characteristics (A—D) and a sagittal view of the copulatory apparatus (E) (after SMITH, 1988; slightly retouched). A, dorsal view of a living specimen; B, anterior end of a living specimen; C, anterior end of a preserved specimen; D, ventral view of the posterior portion of a preserved specimen. Scale, 1 mm. E, sagittal view of the copulatory apparatus of the holotype. Original abbreviations are used (for the new abbreviations shown in parentheses, see explanations in Fig. 4). *a*, atrium (ca); *b*, bursa (cb); *bd*, bursa duct (bs); *co*, common oviduct (cod); *ed*, ejaculatory duct (ma); *gp*, gonopore (gp); *m*, mouth pore (m); *pb*, penis bulb (mz); *ph*, pharynx (ph); *pp*, penis papilla (terminal portion of the mz); *p**v*, papilla of vesicle (pp); *r*, region of ventral gland (ia); *s*, sucker (misunderstanding); *sv*, seminal vesicle (bc); *vd*, vas deference or narrowed spermiductal [sic] vesicle (sd); *vg*, ventral gland (ia).

layers of transverse and circular fibers. Numerous erythrophilic glands lie in the surrounding parenchyma and their outlets penetrate the muscular layer and the inner epithelium (Figs. 4 A—C, 5 A—C; see also Fig. 1 E). A more detailed histology of this organ was found in the original description as "ventral gland" (cf. SMITH, 1988, p. 248, fig. 2, p. 250, fig. 3 b—c).

In histological sections, the moderately large, prepharyngeal testes are arranged in two longitudinal zones, adjacent to each side of the anterior intestinal trunk; they can be seen in preserved specimens (Fig. 2 B and C). They are essentially ventrally located. A pair of ovaries is situated in the usual anteroventral position. Yolk glands are found in dorso-lateral parenchyma on both sides.




Fig. 2. *Seidlia remota* (SMITH, 1988), photographs of two preserved specimens (Lot No. 1985). A and B, dorsal (A) and ventral (B) views of the same specimen; C, ventral view of another specimen. Notice the prepharyngeal testes found in photographs B and C; a pair of spermiducal vesicles can also be seen in the former.

SMITH (1988, p, 248) gave descriptions of anatomy and histology of the copulatory apparatus of "*Pol. remota*" with its schematic figure in sagittal view (*loc. cit.*, fig. 2). The reproduction of his figure is shown in Fig. 1 (E) of the present paper. Our newly prepared sagittal views of the copulatory apparatus of 2 specimens are shown in Fig. 4 (A-C). Pertinent photomicrographs of the apparatus are shown in Fig. 5 (A-C).

The penis consists of a large bulb embedded in the parenchyma and a rather small papilla projecting into the beginning of the male genital antrum. The penis bulb is a napiform organ (Fig. 4 A; sometimes irregular form in shape as shown in Fig. 4 C) and contains a wide bulbar cavity lined by a tall, glandular epithelium. The bulb is pierced many ducts of erythrophilic glands. The two sperm ducts open into the bulbar cavity separately from the antero-lateral sides. The spermiducal vesicles are well-developed in this species (Fig. 4 A and C).

The penis papilla has a short, nearly symmetric and turbinate in shape (Figs. 4 A and C, 5 A and C). A narrow, tubular ejaculatory duct opens at the tip of the papilla. The outer wall of the papilla is lined with a thin epithelium of a nucleate type. The musculature of the papilla is weakly developed. It is noticed in the original description of this species that the structure of the penis papilla and ejaculatory duct were overlooked (*cf.* SMITH, 1988; see Fig. 1 E in the present paper).

The species has an extraordinarily thick muscle zone surrounding the middle and posterior sections of a tubular male antrum that opens into the common genital antrum closely near to the genital pore. The lumen of the male antrum is lined with a cubical, glandular, nucleate epithelium (it is especially well developed in the middle section than other parts). As shown in Fig. 4 (A and C; see also Fig. 5), the posterior end of this muscular organ is bluntly pointed or rounded (so that this part




Fig. 3. *Seidlia remota* (SMITH, 1988), photomicrographs of whole-mounted specimens. Notice the arrangement of eyes. A, head (No. 1985-n; non-stained sample); B, whole body (NO. 1985-o; light stained sample with erythrosin). ia, intestinal anastomosis; gp, genital pore; m, mouth; ph, pharynx.




Fig. 4. *Seidlia remota* (SMITH, 1988), semidiagrammatic sagittal views of the copulatory apparatus of 2 specimens from the type locality. A and B, No. 1985-h (copulatory bursal and bursal stalk are shown separately); C, No. 1985-i
bc, bulbar cavity; bs, bursal stalk; ca, common genital antrum; cb, copulatory bursa; cg, cement glands;



Fig. 5 (continued in the next page).

cod, common ovovitelline duct; ed, ejaculatory duct; gp, genital pore; i, intestine; ia, intestinal anastomosis; m, mouth; ma, male genital antrum; mz, muscle zone surrounding the male genital antrum; od, ovovitelline duct; pb, penis bulb; ph, pharynx; phl, pharynx lumen; pp, penis papilla; sd, sperm duct; sv, spermiducal vesicle; v, vagina.




Fig. 5 (on pp. 101-102). *Seidlia remota* (SMITH, 1988), photomicrographs of near midsagittal sections of the copulatory apparatus. A, No. 1985-h (anterior is the left); B and C, No. 1985-i (anterior is the right). bc, bulbar cavity; bs, bursal stalk; ca, common genital antrum; cb, copulatory bursa; cod, common ovovitelline duct; ed, ejaculatory duct; ia, intestinal anastomosis; m, mouth; ma, male genital antrum; pb, penis bulb; ph, pharynx; phl, pharynx lumen; pp, penis papilla; ps, penis sheath (*i. e.*, muscle zone surrounding the male genital antrum); sd, spermiducal vesicle.

was misunderstood as penis papilla in SMITH's 1988 paper). Histologically, the musculature consists of thin layers of longitudinal fibers and a very thick layer of circular fibers (Figs. 4 A, 5 A and C).

The common ovovitelline duct is rather long and conspicuous. It opens into the terminal portion of the common antrum from the postero-dorsal side (Figs. 4 A and C, 5 B).

The copulatory bursa is moderate in size and is situated in a dorsal position between the posterior end of the pharyngeal chamber and the anterior side of the penis bulb. From the postero-dorsal side of the bursa a wide and long bursal stalk proceeds posteriorly and opens, from the dorsal side, into the terminal portion of the common antrum, immediately near the genital pore (Figs. 4 A-C, 5 A-C). The bursal canal is lined with a tall, glandular,

nucleate epithelium. Below this epithelium, there is a thin layer of circular muscle fibers and a thin layer of longitudinal ones. The posterior one-third of the bursal canal forms a thick-walled vagina. The thick muscle coat surrounding the vagina consists of three layers: an inner, thin layer of longitudinal fibers; a middle, thick layer of circular and longitudinal fibers intermingled; and an outer, slightly thickened layer of longitudinal fibers.

Differential diagnosis. *Seidlia remota* differs from the other members of the genus in the following characters: living animal moderate in size, rather slender, and pale reddish brown above in color; head gently convex with moderately-developed auricles; curved eye-bands separated at the top of the frontal margin and each of them terminating at the anterior one-third level of the prepharyngeal region; possessing a well-developed anastomosis; penis bulb

large and weak muscular with wide bulbar cavity into which each sperm duct enters separately from the antero-lateral sides; turbinate, symmetric penis papilla short and weak muscular with a narrow ejaculatory duct; with an extraordinarily thick muscle zone surrounding the male genital antrum; common ovovitelline duct long which enters into the roof of the posterior terminal part of the common genital antrum; copulatory bursa moderate, with long, rather wide bursal stalk, of which posterior section of the bursal canal forms a thick-walled vagina.

Material. See the section "Material and Methods". Thirteen sets of serial sagittal, transverse and horizontal sections (KAWAKATSU's Specimen Lot No. 1985: -a, -b, -c, -d, -e, -f, -g, -h, -i, -j, -k, -l, -m) and 2 whole mounts (-n, -o) are retained in KAWAKATSU's collection. Some of the sections separated from this collection will be sent to the Department of Zoology (Nat. Hist.), National Science Museum of Natural History, Tôkyô, Japan.

Taxonomic Notes on *Polycelis* and *Seidlia* Species from the Far East and Central Asia

Asia has a rich fauna of the *Polycelis* and *Seidlia* species (cf. KENK, 1953, 1974). Moreover, during the past 10+ years, several new *Polycelis* (s. l.) species were described in this geographical area of Asia. Every known Asiatic species of the genera *Polycelis* and *Seidlia* are listed below with their distribution areas and short taxonomic remarks. For *Polycelis* (*P.*) *sapporo* and *Seidlia schmidtii* from the Far Eastern Area, the latest distribution data (including maps showing their geographical distribution) were described in a recent paper by KAWAKATSU & TIMOSHKIN (In press).

Worldwide distribution of the genera *Polycelis* (subgenera: *Polycelis* and *Ijimia*) and *Seidlia* is shown in Fig. (see KAWAKATSU, OKI, TAMURA, TAKAI, YAMAMOTO, NISHINO, TIMOSHKIN, KUZNEDELOV & SLUYS, 1996: 16, fig. 22).

J A P A N

Polycelis (*Polycelis*) *sapporo* (IJIMA et KABURAKI, 1916). *Olim Sorocelis sapporo*: IJIMA & KABURAKI (1916: 166-167, figs. 19-20); KABURAKI (1922: 32-39, fig. 11, pl. I, fig. 12). Cf. KENK (1953, 1974: 57); HYMAN (1934); OKUGAWA (1938); KAWAKATSU, YANAGITA, TESHIOGI & ICHIKAWA (1974).

Distribution. The lowland, hilly and low mountain districts of Northern Japan (from Aomori Pref. located in the northernmost part of Honshû to the entire region of Hokkaidô (KAWAKATSU & TIMOSHKIN, In press, fig. 1). The species is also recorded from South Sakhalin (PORFIRJEVA & SABITOVA, 1986; PORFIRJEVA, 1990). OKUGAWA's (1953) records in the Middle and South Kuril Islands are uncertain. For other distribution data, see KAWAKATSU & TIMOSHKIN (In press).

Seidlia auriculata (IJIMA et KABURAKI, 1916).

Olim Polycelis auriculata: IJIMA & KABURAKI (1916: 167-169, figs. 21-22); KABURAKI (1922: 39-43, figs. 12-13, pl. I, figs. 13-14). Cf. KENK (1953: 172; 1973: 9, 11-13, figs. 6 D, 9; 1974: 54); KAWAKATSU, TESHIOGI & ISHIDA (1980).

Distribution. Mountain districts of Central and Northern Japan (from central part of Honshû to the Daisetsu Mts. of Hokkaidô) (KAWAKATSU, YAMADA, MURAYAMA & NAOKI, 1991: 356, fig. 1; see also KAWAKATSU, OKI, TAMURA, OGREN, YAMADA & MURAYAMA, 1990: 12, fig. 21); KAWAKATSU (1965a, 1967, 1969, 1974); KAWAKATSU, TESHIOGI & ISHIDA (1980). The record of this species in South Sakhalin (DYGANOVA & PORFIRJEVA, 1990: 59-61, figs. 31-32) is doubtful.

Seidlia akkeshi (ICHIKAWA et KAWAKATSU, 1963). *Olim Polycelis akkeshi*: ICHIKAWA & KAWAKATSU (1963: 2-15, figs. 1-7). Cf. KENK (1974: 54).

Distribution. The southeastern, northeastern and central parts of Hokkaidô (KAWAKATSU, 1969: 74-75, fig. 14, p. 91, pl. VIII, fig. 13).

Seidlia schmidtii (ZABUSOV, 1916). *Olim Rjabu schinskya schmidtii*: ZABUSOV (1916: 273-286, figs. 1-5); *Polycelis ijimai* (KABURAKI, 1917: 327-331, figs. 1-2; see also KABURAKI, 1922: 43-47, fig. 14, pl. I, fig. 15). Cf. ZABUSOVA (1936); LIVANOV & ZABUSOVA (1940); ZABUSOVA-ZHDANOVA (1956, 1960); KENK (1953, 1974, 1989); KAWAKATSU (1964); KAWAKATSU & YAMADA (1966a).

Distribution. The vicinity of Lake Shikotsu-ko, the Lake Akan-ko area, the Shiretoko Peninsula, the Mts. Daisetsu area, the Kitami Mts., the Teshio Mts., and Rishiri Island, in Hokkaidô (KAWAKATSU & TIMOSHKIN, 1998: In press, fig. 2). The species is also recorded from South Sakhalin (ZABUSOVA-ZHDANOVA, 1960; DYGANOVA & PORFIRJEVA, 1990), Primorskiy and Habarovsk in the Russian Far East (PORFIRJEVA & TIMOSHKIN, 1984); the Chukotskiy (Chukotskia) Peninsula (ZABUSOVA-ZHDANOVA, 1960; PORFIRJEVA & TIMOSHKIN, 1980; see also DYGANOVA & PORFIRJEVE, 1990), the Kamchatka Peninsula (ZABUSOVA, 1929, 1936; ZABUSOVA-ZHDANOVA, 1956; PORFIRJEVA, UMYLINA & SHCHEGLOVA, 1979), and Bering Island in the Komandorskiye Islands. MIYADI's (1937) and OKUGAWA's (1953) records of this species from the Kuril Islands could not be confirmed. For other distribution data, see KAWAKATSU & TIMOSHKIN (In press).

SAKHALIN and KAMCHATKA, RUSSIA

Polycelis (*Polycelis*) *elongata* (ZABUSOVA, 1929). *Olim Sorocelis* (*Sorocelides*) *elongata*: ZABUSOVA (1929: 518-521, figs. 8-9, pl. 4, fig. 4). Cf. KENK (1953: 172; 1974: 55). Syn. *Sorocelis relicta* ZABUSOVA, 1929; *Sorocelis eudendrocoeloides* ZABUSOVA, 1929; *Polycelis polyopsis* ZABUSOVA, 1936; *Seidlia eurantron* ZABUSOVA, 1936 (cf. DYGANOVA & PORFIRJEVA, 1990: 64-66, figs. 34-35).

Distribution. Sakhalin and Kamchatka; Bering Island (ZABUSOVA-ZHDANOVA, 1956; PORFIRJEVA & SHCHEGLOVA,

1979; PORFIRJEVA, UMYLINA & SHCHEGLOVA, 1979; DYGANOVA & PORFIRJEVA, 1990). This is a non-pigmented species; the posterior half of the bursal stalk has a thick muscular coat consisting of circular muscle fibers.

? *Polycelis karafto* IJIMA et KABURAKI, 1916. IJIMA & KABURAKI (1916: 169–171, figs. 23–24, pl. I, fig. 16); KABURAKI (1922: 47–49, pl. I, fig. 16). Species inquirenda. Cf. KENK (1953: 172; 1974: 55).

According to the taxonomic descriptions of this species from the vicinities of Yuzhno-Sakhalinsk (Toyohara) and Korsakov (Ôdomari or Korsakoff) in IJIMA & KABURAKI (*loc. cit.*) and KABURAKI (*loc. cit.*), its external appearance is somewhat similar to that of *S. schmidtii*. Although a thick muscle zone of the male genital antrum is not found in the simple schematic figure of the sagittal view of the copulatory apparatus (IJIMA & KABURAKI, 1916: 170, fig. 24), it is highly probable that *Pol. karafto* was named and described from a non-fully matured animal of *S. schmidtii*.

PRIMORSKIY and HABAROVSK, RUSSIA, etc.

Dr. TIMOSHIKUN has a sample (slides) of *Polycelis* sp. from Upper Ussury station (see KAWAKATSU, OKI, TAMURA, YAMAMOTO, NISHINO, TIMOSHIKIN, KUZNEDELOV & SLUYS, 1996: 13, fig. 17); KAWAKATSU has not yet identified this sample. Recently, a few specimens of *Polycelis* sp. were collected by Mr. N. INOUE in a tributary in the vicinity of Habarovsk (Sept. 4, 1997: KAWAKATSU's Specimen Lot No. 2285); *Phagocata* sp. (KSL No. 2284) also occurred. There are no records of identified species of *Polycelis* and/or *Seidlia* in the Habarovsk area and the Korean Peninsula (KAWAKATSU, 1994: 53, fig. 3; KAWAKATSU & LIU, 1987: 44–46, figs. 1–2).

CHINA

Until recently, only 2 described (of which one named only by external appearance of non-sexual specimens) and one non-identified *Polycelis* species were known in China (ZABUSOV, 1911; KATÔ, 1950). In his 1993 book, LIU reported 5 new *Polycelis* species from China (see also LIU, 1994a, b, c, 1995). LIU (1996) also reported 3 additional new *Polycelis* species (of which one was placed in the genus *Seidlia* by KAWAKATSU, 1996) from China. These taxonomic papers cited above were published mainly in Russian, Japanese or Chinese. For Chinese species, KAWAKATSU (1994, 1996) published English corrective papers including reproductions of Chinese descriptions and figures of LIU's (*op. cit.*) 8 species.

Polycelis (Polycelis) wutaishanica LIU, 1993. *Olim Polycelis wutaishanica*: LIU (1993: 160, 162, 166–170, figs. 1–4; 1996: 389–391, figs. 1–2, 397, pl. 1, figs. 1–4). Cf. KAWAKATSU (1994: 53, 55, 58, 60–61, figs. 3, 5–1, 6–1).

Distribution. Shanxi (Shanshi) Province, Northeast-

ern China (LIU, 1993, 1994b, c, 1996; KAWAKATSU, 1994, 1996). *Polycelis* sp. from Shanshi reported by KATÔ (1950: 189, fig. 3) is probably conspecific with this species (KAWAKATSU & LIU, 1987: 45, fig. 1).

Polycelis (Polycelis) jinglensis LIU, 1996. *Olim Polycelis jinglensis*: LIU (1996: 391–393, figs. 3–4, 397, pl. 1, figs. 5–8). Cf. KAWAKATSU (1996: 1–7, figs. 2–1, 3–1, 4).

Distribution. Mt. Tianzhu and Qingchayan, Jingle Counties, Shanxi Province, Northeastern China (LIU, 1996; KAWAKATSU, 1996).

Polycelis (Polycelis) jingyuanica LIU, 1996. *Olim Polycelis jingyuanica*: LIU (1996: 393–394, figs. 5–6, 398, pl. II, figs. 1–4). Cf. KAWAKATSU (1996: 1–7, figs. 2–2, 3–2, 4).

Distribution. Laolongtan and Xixia Jingyuan Counties, Ningxia Huiz Autonomous Region, Central China (LIU, 1996; KAWAKATSU, 1996).

Seidlia hamica (LIU, 1996). *Olim Polycelis hamica*: LIU (1996: 394–396, figs. 7–8, 398, pl. II, figs. 5–8). Cf. KAWAKATSU (1996: 1–7, figs. 2–3, 3–3, 4).

Distribution. Baishan of Qincheng, Hami County, Xinjiang Uygur Autonomous Region, Northwestern China (LIU, 1996; KAWAKATSU, 1996).

Polycelis (Polycelis) koslowi (ZABUSOV, 1911). *Olim Sorocelis koslowi*: ZABUSOV (1911: 345–348, pls. II, fig. 4, XI, fig. 3). Cf. KENK (1953: 172; 1974: 55).

Distribution. Xizang Autonomous Region (Tibet), Southwestern China (ZABUSOV, 1911; PORFIRJEVA, 1960: 122–131, figs. 1–2; DYGANOVA & PORFIRJEVA, 1990: 72–74, fig. 43; KAWAKATSU, 1994: 53, 64, figs. 3, 7).

Sorocelis tibeticva ZABUSOV, 1911, is a species inquirenda reported from Tibet; its genital anatomy is not known (ZABUSOV, 1911: 349–350, pl. II, fig. 1; see KENK, 1974: 57; KAWAKATSU, 1994: 64, 67, fig. 7). Judging from the external morphology of this species in the original description, it is highly probable that *Sor. tibetica* [nec *Polycelis tibetica* HYMAN, 1934] is a synonym of *P. (P.) koslowi*. *Polycelis* sp. from Nepal (ICHIKAWA & KAWAKATSU, 1964: 27–29, figs. 1–3) may be a synonym of *P. (P.) koslowi*.

Polycelis (Polycelis) sinensis LIU, 1993. *Olim Polycelis sinensis*: LIU (1993: 154, 156, 160, 162–163, fig., 172–174, fig. 1). Cf. KAWAKATSU (1994: 53, 55–56, 58, 60–61, 64, figs. 5–2, 6–2, 7).

Distribution. Xizang Autonomous Region, Southwestern China (LIU, 1993, 1994b, c, 1995; KAWAKATSU, 1994).

Polycelis (Polycelis) lhunzhubica LIU, 1993. *Olim Polycelis lhunzhubica*: LIU (1993: 160, 162–163, fig., 174–175, fig. 2). Cf. KAWAKATSU (1994: 53, 56, 59–60, 62, figs. 3, 5–3, 6–3, 7).

Distribution. Xizang Autonomous Region, Southwestern China (LIU, 1993, 1994b, c, 1995; KAWAKATSU, 1994).

Polycelis (Polycelis) nyingchica LIU, 1993. *Olim Polycelis nyingchica*: LIU (1993, 161–63, fig., 176–178, figs. 1–2). Cf. KAWAKATSU (1994: 53, 56, 59–60,




Fig. 6. Sagittal views of the copulatory apparatus of *Polycelis* (*Polycelis*) *pathan* DE BEAUCHAMP, 1959. A (top), *Polycelis pathan* from Afghanistan (after DE BEAUCHAMP, 1959); B (middle and bottom), "*Polycelis asiatica*" from Tadzhikistan (after SELIVANOVA, 1985). Only French abbreviations are given. b, bourse copulatrice; cd, canal déferent; gb, glandes de la bourse; gc, glandes de l'atrium; mb, musculature bulbaire; p, pénis; vs, vésicule séminale.

64, figs. 3, 5-4, 6-4, 7).

Distribution. Xizang Autonomous Region, South-western China (LIU, 1993, 1994a, b, c; KAWAKATSU, 1994).

Polycelis (Polycelis) xigazensis LIU, 1993. *Olim polycelis xigazensis*: LIU (1993, 161-163, fig., 176-180, figs. 3-4). Cf. KAWAKATSU (1994: 53, 56, 59-60, 63-64, figs. 3, 5-5, 6-5, 7).

Distribution. Xizang Autonomous Region, South-western China (LIU, 1993, 1994a, b, c, KAWAKATSU, 1994).

Mr. LIU's 4 *Polycelis* species reported from Tibet show striking resemblances to one another. The genital anatomy and histology of *Pol. (P.) sinensis* are the same as those of *Pol. (P.) lhunzhubica*. Moreover, *Pol. (P.) nyingchica* is probably a synonym of *Pol. (P.) koslowi*. These problems will await future study.

Polycelis (Polycelis) kashimirica LIU, 1993. *Olim Polycelis tibetica* HYMAN, 1934 [nec *Sorocelis tibetica* ZABUSOV, 1911]. Cf. HYMAN (1934: 7-11, pls. I, figs. 1-8, II figs. 1-5); KENK (1953: 172; 1974: 57); LIU (1993: 181-182); KAWAKATSU (1994: 53, 64-66, figs. 3, 7).

Distribution. Kashmir (Indian Tibet) (HYMAN, 1934: KAWAKATSU, 1994). The genital anatomy and histology of this species have great resemblance to those of *Pol. (P.) koslowi* from Tibet.

NEPAL

Polycelis sp. (spp. ?) and/or *Seidlina* sp. (spp. ?) of Nepal. Non-identified samples in KAWAKATSU's Collection.

During the past 35+ years, KAWAKATSU obtained a considerable number of preserved specimens of *Dugesia* sp. (spp. ?) and *Polycelis* sp. (spp. ?) and/or *Seidlina* sp. (spp. ?) collected by many friends in the Kingdom of Nepal (cf. KAWAKATSU, YONEDA & MURAYAMA, 1996a: 95-103, fig. 1; 1996b: 1-4, figs. 1-7) (alt. 1525-4300 m). Several additional samples from the Khumbu Himal area (alt. 2440-4755 m) were also received from Dr. A. OHTAKA. All of these samples sent to Dr. R. SLUYS will be studied with him.

TADZHIKISTAN; KAZAKHSTAN and AFGHANISTAN (CENTRAL ASIA)

Polyselis (Polycelis) pathan DE BEAUCHAMP, 1959. *Olim Polycelis pathan*: DE BEAUCHAMP (1959: 37-42, figs. 6-8). Cf. KENK (1974: 56). Syn. *Polycelis asiatica* SELIVANOVA, 1985. syn. nov. Cf. SELIVANOVA (1985: 820-826, figs. 1-2). (For comparison of the sagittal views of the copulatory apparatus of this species, see Fig. 6A and




Fig. 7. Worldwide distribution of the genera *Polycelis* EHRENBERG, 1831 (subgenera *Polycelis* EHRENBERG, 1831, and *Ijimia* BERGENDAL, 1890) and *Seidlina* ZABUSOV, 1911. (After KAWAKATSU, OKI, TAMURA, TAKAI, YAMAMOTO, NISHINO, TIMOSHIN, KUZNEDELOV & SLUYS, 1996; modified.)

B in the present paper.)

Distribution. Afghanistan and Tadzhikistan (Tadjikistan) (KAWAKATSU, 1973 : 80, fig. 1; DYGANOVA & PORFIRJEVA, 1990 : 74-75, fig. 44).

Polycelis (Polycelis) pamirensis DE BEAUCHAMP, 1961. *Olim Polycelis pamirensis* : DE BEAUCHAMP (1961 : 78-85, figs. 1-3); ICHIKAWA & KAWAKATSU (1966 : 176-179, 182-187, figs. 1-6). Cf. KENK (1974 : 56).

Distribution. Afghanistan (ICHIKAWA & KAWAKATSU, 1966; KAWAKATSU, 1973 : 80, fig. 1). *Polycelis* sp. from Northern Pakistan (KAWAKATSU, 1973 : 82-83, figs. 2 C-E, 3 D, 88-89) is probably conspecific with this species.

Seidlia sabussowi (SEIDL, 1911). *Olim Sorocelis sabussowi* : SEIDL (1911 : 36-64, pls. V-VIII, figs. 6, 11-12). Cf. KENK (1953 : 172; 1974 : 56-57). Syn. *Sorocelis stummeri* SEIDL, 1911; *Sorocelis lactea* SEIDL, 1911; *Seidlia alma-atina* ZABUSOVA-ZHDANOVA, 1947. Cf. SEIDL (1911); ZABUSOV (1911); ZABUSOVA-ZHDANOVA (1947 : 153-160, figs. 1-3); DYGANOVA & PORFIRJEVA (1990 : 62-63, fig. 33).

Distribution. Karakul' in Tadzhikistan and Kazakhstan.

Polycelis (Polycelis) kulsai ZABUSOVA-ZHDANOVA, 1947. *Olim Polycelis kulsai* : ZABUSOVA-ZHDANOVA (1947 : 160-164, figs. 4-6). Cf. KENK (1974 : 55).

Distribution. Kazakhstan (PORFIRJEVA, 1960 : 131-138, figs. 3-4).

KAVKAZ, RUSSIA (CENTRAL ASIA)

Polycelis (Polycelis) eburnea (MUTH, 1912). *Olim Sorocelis eburnea* : MUTH (1912 : 387-408, figs. 1-2). Cf. KENK (1953 : 172; 1974 : 54-55).

Distribution. Kavkaz, Russia (PORFIRJEVA, 1960; DYGANOVA & PORFIRJEVA, 1990 : 70-72, fig. 41).

ACKNOWLEDGEMENTS

For the supply of specimens of *Seidlia remota* collected from the type locality, the authors wish to thank Dr. Douglas G. SMITH (Amherst, Massachusetts, U. S. A.). They are also grateful to the following gentlemen for discussing the taxonomic problem on *Polycelis* and *Seidlia* groups : Dr. Oleg A. TIMOSHIN (Irkutsk, Russia), Dr. Robert E. OGREN (Kingston, Pennsylvania, U. S. A.) and Dr. Ronald SLUYS (Amsterdam, The Netherlands). Finally, KAWAKATSU is indebted to the following Japanese gentlemen for sending him recent collection of freshwater planarians from Habarovsk in Russia and Nepal : Mr. Nobuo INOUE (Nigata), Mr. Hitoshi MURAYAMA (Nagaoka) and Dr. Akifumi OHTAKA (Hiroaki).

SUMMARY

"*Polycelis remota* SMITH, 1988," described from a spring in western Massachusetts, is a medium-sized, pig-

mented species. From the results of the microscopic examination of newly-prepared serial sections of animals from the type locality, the species has a small penis papilla projecting into the beginning of the long, tubular male genital antrum that is surrounded by a thick, highly muscular coat (penis sheath). The species should be classified as *Seidlia remota* (SMITH, 1988).

The taxonomic notes on *Seidlia* and *Polycelis* species from the Far East and Central Asia were given in the second section. "*Polycelis asiatica* SELIVANOVA, 1985," from Tadzhikistan should be considered as a synonym of *Polycelis (Polycelis) pathan* DE BEAUCHAMP, 1959, from Afghanistan.

REFERENCES

- BEAUCHAMP, P. DE, 1959. Triclaides paludicoles d'Afghanistan (Contribution à l'étude de la faune d'Afghanistan 1). Kungl. Fysiogr. Aälsk. I Lund Förhandl., 29 (3) : 27-43. 1961. Triclaides Paludicoles d'Afghanistan (deuxième note). *Ibid.*, 31 (6) : 77-85.
- BERGENDAL, D., 1890. Studien über nordische Turbellarien und Nemertinen. Öfver. Kongl. Vetensk.-Akad. Förhandl., 47 : 323-328.
- DYGANOVA, R. YA. & PORFIRJEVA, N. A., 1990. *Planarii Aziatskoï Chasti SSSR*. Morfologiya Sistematika, Rasprostranenie. Pp. 1-151. Izdatel'stvo Kazanskogo Univ., Kazan.
- EHRENBERG, C. G., 1831. Animalium evertibrata exclusis Insectis. In: HEMPRICH, F. G. & EHRENBERG, C. G. (eds.), "Symbolae Physicae". Vol. 4, series prima cum tabularum decade prima, Berolini.
- HYMAN, L. H., 1934. Report on Triclad Turbellaria from Indian Tibet. Mem. Connecticut Acad. Arts & Sci., 10 (2) : 5-12 + pls. I-II.
- ICHIKAWA, A. & KAWAKATSU, M., 1963. *Polycelis akkeshi*, a new freshwater planarian, from Hokkaido. Publ. Akkeshi Mar. Biol. Stat., (12) : 1-18. 1964. Report on freshwater planaria from Northeast Nepal. Zool. Mag., Tokyo, 73 : 27-31. (In Japanese, with English summary.) 1966. Report on freshwater planaria from Afghanistan. In : KITAMURA, S. & YOSII, R. (eds.), Results of the Kyoto Univ. Sci. Exped. to the Karakoram and Hindukush, 1955, VIII, Additional Reports. Pp. 175-187. Comm. Kyoto Univ. Sci. Exped. etc., Kyôto Univ., Kyôto.
- IJIMA, I. & KABURAKI, T., 1916. Preliminary descriptions of some Japanese triclads. Annot. Zool. Japon., 9 : 153-171.
- KABURAKI, T., 1917. Notizen über japanische Tricladen. Annot. Zool. Japon., 9 : 325-333. 1922. On some Japanese freshwater triclads ; with a note on the parallelism in their distribution in Europe and Japan. Jour. Coll. Sci. Imp. Univ. Tokyo, 44 : 1-71 + pl. I.
- KATÔ, K., 1950. On some Turbellarians from Sanshi, North China. Zool. Mag., Tokyo, 59 : 188-190. (In Japanese.)

- KAWAKATSU, M., 1964. Studies on the taxonomy and morphology of the Japanese freshwater planarian, *Polycelis schmidtii* (ZABUOV). Annot. Zool. Japon., 37: 174–184. 1965. On the ecology and distribution of freshwater planarians in the Japanese Islands, with special reference to their vertical distribution. Hydrobiologia, 26: 349–408. 1967. On the ecology and distribution of freshwater planarians in the Japanese Islands, with special reference to their vertical distribution (Revised edition). Bull. Fuji Women's College, (5): 117–177. 1969. An illustrated list of Japanese freshwater planarians in color. *Ibid.*, (7), II: 45–91 (+ pls. VII–VIII). 1973. Report on freshwater planarians from Pakistan. *Ibid.*, (11), II: 79–95. 1974. Further studies on the vertical distribution of freshwater planarians in the Japanese Islands. In: RISER, N. W. & MORSE, M. P. (eds.), *The HYMAN Memorial Volume ... Biology of the Turbellaria*. Pp. 291–338. McGraw-Hill Book Co., New York, etc. 1994. A commentary note on Prof. LIU's 1993 and 1994 Chinese Publications on Freshwater planarians from China, with corrections of the date of original publications of five *Polycelis* species and a replacement name of "*Polycelis tibetica* HYMAN, 1934". Bull. Fuji Women's College, (32), II: 45–71. 1996. A commentary note on Prof. LIU's 1996 Chinese publication on freshwater planarians from China. Occ. Publ., Biol. Lab. Fuji Women's College, Sapporo (Hokkaidô), Japan, (29): 1–8.
- KAWAKATSU, M. & LIU, D.-Z., 1987. History of the study of Turbellaria in China. Part 3. Supplementary notes on the turbellariology in the People's Republic of China. Bull. Fuji Women's College, (25), II: 39–54.
- KAWAKATSU, M. & MITCHELL, R. W., 1995. New taxonomic data on an American triclad, *Seidlia remota* (SMITH, 1988), with a proposal that generic recognition be extended to the subgenus *Seidlia*. Zool. Sci., Tôkyô, 12-Suppl.: 32.
- KAWAKATSU, M., OKI, I., TAMURA, S., OGREN, R. E., YAMADA, T. & MURAYAMA, H., 1990. Preprint of papers given at the Sixth International Symposium on the Turbellaria, Hirosaki, Japan, Augst 7–12, 1990. Occ. Publ., Biol. Lab. Fuji Women's College, Sapporo (Hokkaidô), Japan (22): 1–16.
- KAWAKATSU, M., OKI, I., TAMURA, S., TAKAI, M., YAMAMOTO, K., NISHINO, M., TIMOSHKIN, O. A., KUZNEDELOV, K. D. & SLUYS, R., 1996. Preprint of papers given at the Eighth International Symposium on the Biology of Turbellaria, Brisbane, Australia, August 19–23, 1996. Occ. Publ., Biol. Lab. Fuji Women's College, Sapporo (Hokkaidô), Japan, (27): 1–16.
- KAWAKATSU, M., TESHIOGI, W. & ISHIDA, S., 1980. Morphological variation of the copulatory apparatus of a freshwater planarian, *Polycelis (Seidlia) auriculata* IJIMA et KABURAKI, collected from five localities in the Tôhoku Region, Honshû, Japan. Sci. Rep. Hirosaki Univ., 27 (2): 130–146.
- KAWAKATSU, M. & TIMOSHKIN, O. A., 1996. The geographic distribution of *Polycelis (Polycelis) sapporo* and *Seidlia schmidtii* in the Far East. VIII ISBT Program/Abstracts (Bribane, August 18–23, 1996): 55. In press. The geographical distribution of *Polycelis (Polycelis) sapporo* and *Seidlia schmidtii* in the Far East. Hydrobiologia.
- KAWAKATSU, M. & YAMADA, T., 1966. Additional studies on the morphology of *Polycelis schmidtii* (ZABUOV), with remarks on distribution. Bull. Biolgeogr. Soc. Japan, 24: 1–8.
- KAWAKATSU, M., T., YAMADA, T., MURAYAMA, H. & NAOKI, Y., 1991. Geographical distribution of *Polycelis (Seidlia) auriculata* in Japan. Hydrobiologia, 227: 355–363.
- KAWAKATSU, M., YANAGITA, Y., TESHIOGI, W. & ICHIKAWA, A., 1974. Studies on the taxonomy and morphology of the Japanese freshwater planarian, *Polycelis sapporo* (IJIMA et KABURAKI, 1916). Annot. Zool. Japon., 47: 239–260.
- KAWAKATSU, M., YONEDA, I. & MURAYAMA, H., 1996a. A list of Triclad Turbellarians from Nepal in KAWAKATSU's Collection. Part I: A list with maps. Bull. Fuji Women's College, (34), II: 95–103. 1996b. Do. Part II: Additional maps and photographs. Occ. Publ., Biol. Lab. Fuji Women's College, Sapporo (Hokkaidô). Japan, (28): 1–4.
- KENK, R., 1930. Beiträge zum System der Probursalier (Tricladida Paludicola). Zool. Anz., 89: 145–162, 289–302. 1953. The fresh-water triclads (Turbellaria) of Alaska. Proc. U. S. Nat. Mus., 103 (3322): 163–186 + pls. 6–8. 1972. *Freshwater Planarians (Turbellaria) of North America*. Biota of Freshwater Ecosystems, Identification Manual. No. 1: i–ix + 1–81. U. S. Environment. Protect Agency, Washington, D. C. 1973. Freshwater Triclad (Turbellaria) of North America, V: The genus *Polycelis*. Smithsonian Contrb. Zool., (135): 1–15. 1974. Index to the genera and species of the freshwater triclads (Turbellaria) of the world. *Ibid.*, (183): i–ii, 1–90. 1989. Revised list of the North American freshwater planarians (Platyhelminthes: Tricladida: Paludicola). *Ibid.*, (476): i–iv, 1–10.
- LIU, D.-Z., 1993. *Chinese Freshwater Turbellarians*. Ten prefatory pages + pp. 1–184. National Peking Normal Univ. Press, Peking. (In Chinese.) 1994a. Notes on two new species of the genus *Polycelis* from Xizang Autonomous Region, China (Tricladida: Paludicola: Planariidae). Acta Zootaxon. Sinica, 19 (1): 10–15 + pl. I. (In Chinese, with English summary.) 1994b. Recent progress of taxonomic study of the genus *Polycelis* (Tricladida) in China. In: Abstracts of the Commemor. Meet. of the Sixtieth Ann. of the China Zool. Soc., p. 118. Peking. (In Chinese.) 1994c. On the taxonomy of Chinese Paludicola (Turbellaria). In: Proceed. of the Sixtieth Ann. of the Found. of China Zool. Soc., 1934–1994, pp. 543–548 + fig. 1. Zhongguo Kexue-Jiguh Publ., Peking. (In Chinese.) 1995. Notes on two new species of genus *Polycelis* from China (Tricladida: Paludicola: Planariidae). Acta Hydrobiol. Sinica, 19 (2): 152–156. (In Chinese, with English summary.) 1996. Notes on four new species of the genus *Polycelis*.

celis from China (Tricladida : Paludicola : Planariidae). Acta Zootaxon. Sinica, 21 (4) : 389-398 + pls. I-II. (In Chinese, with English summary.)

LIVANOV, N. A. & ZABUSOVA, Z. I., 1940. Planariï basseina Teletskogo ozera i novye o nekotorykh drugikh sibirskikh vidakh. Trudy Obshch. Estestvoisp. pri Kazanskom Univ., 56 (3/4) : 83-159. (In Russian, with French summary.)

MIYADI, D., 1937. Limnological survey of the North Kuril Islands. Arch. Hydrobiol., 31 : 433-483 + pls. VII-IX.

MUTH, A., 1912. Beiträge zur Kenntnis der Gattung *Sorocelis* GRUBE. Mitteil. Naturwiss. Vereins Steiermark, 48 : 381-410.

OKUGAWA, K. I., 1938. Probursalia of Hokkaidô, with special reference to *Sorocelis sapporo*. Zool. Mag., Tôkyô, 50 : 177. (In Japanese.) 1953. A monograph of Turbellaria (Acoela, Rhabdocoela, Alloecoela and Tricladida) of Japan and its adjacent regions. Bull. Kyoto Gakugei Univ., B (3) : 20-43.

PORFIRJEVA, N. A., 1960. Planariï Sredonei Azii. Trudy Obshch. Estestvoisp. pri Kazanskom Gosud. Univ., 63 : 122-138. (In Russian.)

PORFIRJEVA, N. A. & SABITOVA, S. E., 1986. Novye dannye o planariakh Iuzhnogo Sakhalina. In: Donnye Organizmy Presnykh vod Dal'nego Vostoka. Akad. Nauk SSSR, Vladivostok : 15-18. (In Russian.)

PORFIRJEVA, N. A. & SHCHEGLOVA, E. G., 1979. Planariï (Tricladida, Paludicola) Ostrova Beringa (Komandorskie Ostrova). Zool. Zhur., Moscow, 58 (1) : 117-119. (In Russian, with English summary.)

PORFIRJEVA, N. A. & TIMOSHKIN, O. A., 1980. Planariï (Tricladida, Paludicola) Chukotskogo poluostrova i ikh zoogeograficheskie svyazi. In: Fauna Presnykh vod Dal'nego Vostoka. Akad. Nauk SSSR, Vladivostok : 37-43. (In Russian.) 1984. K faune planariï (Tricladida, Paludicola) Sikhote-Alinia. In: Biologiya Presnykh vod Dal'nego Vostoka. Akad. Nauk SSSR, Vladivostok : 62-68. (In Russian.)

PORFIRJEVA, N. A., UMYLINA, T. M. & SHCHEGLOVA, E. G., 1979. K revizii founy presnovod-

nykh Triklad (Turbellaria, Tricladida) Kamchatki. In: Sistematika i Ekologiya ryb kontinental'nykh vodemov Dal'nego Vostoka. Akad. Nauk SSSR, Vladivostok : 51-61. (In Russian.)

SEIDL, H. H., 1911. Beiträge zur Kenntnis centralasiatischer Tricladen. Zeitschr. Wiss. Zool., 98 : 31-67 + pls. V-VII.

SELIVANOVA, R. V., 1985. Novyi predstavitel' planariï iz Tadzhikistana (Tricladida, Paludicola). Zool. Zhur., Moscow, 64 (6) : 820-826. (In Russian, with English summary.)

SMITH, D. G., 1988. A new, disjunct species of triclad flatworm (Turbellaria : Tricladida) from spring in southern New England. Biol. Bull., 175 : 246-252.

STIMPSON, W., 1857. Prodrum descriptionis animalium evertibratorum quae in expeditione ad Oceanum Pacificum septentrionalem a Republica Federata missa, Johanne RODGERS duce, observavit et descripsit, I : Turbellaria Dendrocoela. Proc. Acad. Nat. Sci. Philad., 9 : 19-31.

ZABUSOVA, I. P. (=SABUSSOW, H.; ZABOUSSOFF, H.), 1911. Izsledovaniia po morfologii i sistematike planariï ozera Balkala, I : Rod *Sorocelis* GRUBE. Trudy Obshch. Estestvoisp. pri Imp. Kazanskom Univ., 43 (4) : 1-422 + 1-8 + 1-2 + pls. I-XI. (In Russian, with German summary.) 1916. *Rjabuschinskya schmidtii* n. g. n. sp., novyi vid i rod Tricladida paludicola iz' Kamchatki. Rus. Zool. Zhur., Saratov, 1 : 273-286. (In Russian, with French summary.)

ZABUSOVA, Z. I. (=SABUSSOFF, Z.; = SABUSSOWA, Z.; ZABUSOVA-ZHDANOVA, Z. I.), 1929. Die Turbellarien der Kamtschatka-Halbinsel nach den Sammlungen der Rjabuschinsky-Expedition 1908-1909. Zool. Jb. Syst., 57 : 497-356 + pl. IV. 1936. Planariï Kamchatki. Trudy Obshch. Estestvoisp. Kazanskom Univ., 96 (7) : 141-174. (In Russian, with French summary.) 1947. Novye dannye po Sredne-Aziatskim planariam., *Ibid.*, 57 (3/4) : 153-161. (In Russian.) 1956. Planariï nerestilische lososevykh ryb zapadnogo Poberezhia Kamchatki. *Ibid.*, 96 (7) : 25-39. (In Russian.) 1960. Planariï dal'nego Vostoka. *Ibid.*, 63 : 112-138. (In Russian.)

Addresses of the Authors :

Dr. Masaharu KAWAKATSU, Professor of Biology, Fuji Women's College, Kita-16, Nishi-2, Kita-ku, Sapporo (Hokkaidô) 001 - 0016, Japan.

Home address for correspondence (after January, 1999) : 9 jô 9 chôme 1-8, Shinkotoni, Kita-ku, Sapporo (Hokkaidô) 001 - 0909, Japan.

Fax & Tel. (International : +81 - 11 - 762 - 4450) ; (Domestic : 011 - 762 - 4450).

Dr. Robert W. MITCHELL, Professor of Biology, Emeritus, Texas Tech University, Lubbock, Texas, U. S. A.

Home address for correspondence : 21 Windmill Drive, Route 3, Box 921, Bandera, Texas 78003, U. S. A.

Fax (International : +1 - 210 - 796 - 4243).

December 25, 1998.